FARG‘ONA DAVLAT UNIVERSITETI


O‘ZBEKISTON RESPUBLIKASI
OLIY TA‘LIM, FAN VA INNOVATSIYALAR VAZIRLIGI
FARG‘ONA DAVLAT UNIVERSITETI

YANGI AVLOD
ILMIY TADQIQOTLARI

Respublika ilmiy-amaliy konferensiya to‘plami

SHO‘BA:

Aniq fanlar



ILM-FAN YUTUQLARI – YUKSALISH SARI

INNOVATSIYA – KELAJAK POYDEVORI

2026-YIL 25-IYUN

FARG‘ONA

THE TWO-PARAMETRIC MITTAG-LEFFLER FUNCTION AND ITS SOME PROPERTIES

Furkatjon F. Nematjonov,
an assistant at Fergana State Technical University
fnematjonov31@gamil.com

Abstract
In this article it is presented that the basic properties of the two-parametric Mittag-Leffler function $E_{\alpha,\beta}(z)$*, which is the most straightforward generalization of the classical Mittag-Leffler function* $E_{\alpha}(z)$*. Its order and type are calculated, this article presents a number of formulas relating to the two-parametric Mittag-Leffler function's recurrence relations and differentiation, introduce some useful integral representations.*

Keywords: *Gamma function, complementary error function, Prabhakar function, Abel integral equation, Beta function.*

1. Series representation

The Mittag-Leffler type function (or the *two-parametric Mittag-Leffler function*)

$$E_{\alpha,\beta}(z)=\sum_{k=0}^{\infty}\frac{z^k}{\Gamma(\alpha k+\beta)}\ (\mathrm{Re}\,\alpha>0,\ \beta\in\mathbb{C}) \tag{1}$$

Generalizes the classical Mittag-Leffler function

$$E_{\alpha}(z)=\sum_{k=0}^{\infty}\frac{z^k}{\Gamma(\alpha k+1)}\ (\mathrm{Re}\,\alpha>0).$$

We have $E_{\alpha,1}(z)=E_{\alpha}(z)$. For any $\alpha,\beta\in\mathbb{C}$, $\mathrm{Re}\,\alpha>0$, the function (1) is an entire function of order $\rho=1/(\mathrm{Re}\,\alpha)$ and type $\sigma=1$.

Indeed, let us consider the slightly more general function

$$E_{\alpha}(\sigma^{\alpha}z)=\sum_{k=0}^{\infty}\frac{(\sigma^{\alpha}z)^k}{\Gamma(\alpha k+1)}, \tag{2}$$

i.e. take the coefficients in the form

$$c_k=\frac{\sigma^{\alpha k}}{\Gamma(\alpha k+\beta)}\ (k=0,1,2,\ldots) \tag{3}$$

Where $0<\mathrm{Re}\,\alpha<+\infty,\ 0<\sigma<+\infty$ is an arbitrary real constant and β is a complex parameter. By using Stirling's formula we have

$$\Gamma(\alpha k+\beta)=\sqrt{2\pi}(\alpha k)^{\alpha k+\beta-\frac{1}{2}}e^{-\alpha k}\left[1+o(1)\right],\ k\to\infty \tag{4}$$

And, consequently, for the sequence $\{c_k\}_0^{\infty}$ we immediately obtain

$$\lim_{k\to\infty}\frac{k\log k}{\log\frac{1}{|c_k|}}=\frac{1}{(\mathrm{Re}\,\alpha)}=\rho,\quad \lim_{k\to\infty}k|c_k|^{\rho/k}=e\rho\sigma \tag{5}$$

2. Differential and Recurrence Relations

A term-by-term differentiation allows us to verify in an easy way that

$$\frac{d^m}{dz^m}E_{\alpha,\beta}(z)=\sum_{k=0}^{\infty}\frac{\Gamma(k+m+1)z^k}{k!\Gamma(\alpha k+\alpha m+\beta)}=m!E^{m+1}_{\alpha,\alpha m+\beta}(z)\quad(m\geq 1) \tag{6}$$

Where $E^{\gamma}_{\alpha,\beta}(z)$ denotes the 3-parametric Mittag-Leffler function (also known as the Prabhakar function).

The following formula expresses the first derivative of the ML function in terms of the difference of two instances of the same function

$$\frac{d}{dz}E_{\alpha,\beta}(z)=\frac{E_{\alpha,\beta-1}(z)+(1-\beta)E_{\alpha,\beta}(z)}{\alpha z},\ z\neq 0 \tag{7}$$

It can be generalized to derivatives of any integer order, and thus provides a summation formula of Djrbashian type: *Let* $\alpha>0,\beta\in$ □ *and* $z\neq 0$. For any $m\in$ □

$$\frac{d^m}{dz^m}E_{\alpha,\beta}(z)=\frac{1}{\alpha^m z^m}\sum_{j=0}^{m}c_j^{(m)}E_{\alpha,\beta-j}(z) \tag{8}$$

where $c_0^{(0)}=1$ *and the remaining coefficients* $c_j^{(m)}$, $j=0,1,...,m$, *are recursively evaluated as*

$$c_j^{(m)}=\begin{cases}(1-\beta-\alpha(m-1))c_0^{(m-1)} & j=0,\\ c_{j-1}^{(m-1)}+(1-\beta-\alpha(m-1)+j)c_j^{(m-1)} & 1\leq j\leq m-1,\\ 1 & j=m\end{cases} \tag{9}$$

Several applications involve the more general function $z^{\beta-1}E_{\alpha,\beta}(z^{\alpha})$, for which differentiation formulas can easily be derived. The following differentiation formula is an immediate consequence of the definition of the two-parametric Mittag-Leffler function (1)

$$\left(\frac{d}{dz}\right)^m\left[z^{\beta-1}E_{\alpha,\beta}(z^{\alpha})\right]=z^{\beta-m-1}E_{\alpha,\beta-m}(z^{\alpha})\quad(m\geq 1) \tag{10}$$

We consider now some corollaries of formula (10). Let $\alpha=m/n,(m,n=1,2,...)$ in (10). Then

$$\left(\frac{d}{dz}\right)^m\left[z^{\beta-1}E_{m/n,\beta}\left(z^{m/n}\right)\right]=$$

$$=z^{\beta-1}E_{m/n,\beta}\left(z^{m/n}\right)+z^{\beta-1}\sum_{k=1}^{n}\frac{z^{-\frac{m}{n}k}}{\Gamma\left(\beta-\frac{m}{n}k\right)}\quad(m,n\geq 1) \tag{11}$$

Since

$$\frac{1}{\Gamma(-s)}=0\quad(s=0,1,2,\ldots),$$

It follows from (11) with n=1 and $\beta=0,1,\ldots,m$ that

$$\left(\frac{d}{dz}\right)^m\left[z^{\beta-1}E_{m,\beta}\left(z^m\right)\right]=z^{\beta-1}E_{m,\beta}\left(z^m\right)\quad(m\geq 1) \tag{12}$$

Substituting $z^{n/m}$ in place of z in (11) we get

$$\left(\frac{m}{n}z^{1-\frac{n}{m}}\frac{d}{dz}\right)^m\left[z^{(\beta-1)\frac{n}{m}}E_{m,\beta}(z)\right]$$

$$=z^{(\beta-1)\frac{n}{m}}E_{m,\beta}(z)+z^{(\beta-1)\frac{n}{m}}\sum_{k=1}^{n}\frac{z^{-k}}{\Gamma\left(\beta-\frac{m}{n}k\right)}\qquad(m,n=1,2,\ldots) \tag{13}$$

Let $m=1$ in this formula. We then obtain the first-order differential equation for the function $z^{(\beta-1)n}E_{1/n,\beta}(z)$:

$$\frac{1}{n}\frac{d}{dz}\left[z^{(\beta-1)n}E_{1/n,\beta}(z)\right]-z^{n-1}\left[z^{(\beta-1)n}E_{1/n,\beta}(z)\right]$$

$$=z^{\beta n-1}\sum_{k=1}^{n}\frac{z^{-k}}{\Gamma\left(\beta-\frac{k}{n}\right)},\quad(n=1,2,\ldots) \tag{14}$$

Here we consider $u=z^{(\beta-1)n}E_{1/n,\beta}(z)$; $f(z)=z^{\beta n-1}\sum_{k=1}^{n}\frac{z^{-k}}{\Gamma\left(\beta-\frac{k}{n}\right)}$ in order to solve the non-homogeneous ordinary differential equation, and then

$$\frac{1}{n}\frac{du}{dz}-z^{n-1}u=f(z)$$

Solving this equation, we obtain for any $z_0\neq 0$,

$$E_{1/n,\beta}(z)=z^{(1-\beta)n}e^{z^n}\left\{z_0^{(\beta-1)n}e^{-z_0^n}E_{1/n,\beta}(z_0)+n\int_{z_0}^{z}e^{-\tau^n}\left(\sum_{k=1}^{n}\frac{\tau^{-k}}{\Gamma\left(\beta-\frac{k}{n}\right)}\tau^{\beta n-1}\right)d\tau\right\}\quad(n=1,2,\ldots)$$

(15)

Formula (15) is true with $z_0=0$ if $\beta=1$. In this case we have

$$E_{1/n,1}(z)=e^{z^n}\left\{1+n\int_0^z e^{-\tau^n}\left(\sum_{k=1}^{n-1}\frac{\tau^{k-1}}{\Gamma\left(\frac{k}{n}\right)}\right)d\tau\right\}(n\geq 2)\qquad(16)$$

In particular,

$$E_{1/2,1}(z)=e^{z^2}\left\{1+\frac{2}{\sqrt{\pi}}\int_0^z e^{-\tau^2}d\tau\right\}=e^{z^2}\{1+\operatorname{erf}z\}=e^{z^2}\operatorname{erfc}(-z)\qquad(17)$$

And, consequently,

$$E_{1/2,1}(z)\square\ 2e^{z^2},\ |\arg z|<\frac{\pi}{4},\ |z|\to\infty$$

In the following lemma we collect together a number of known recurrence relations for the two-parametric Mittag-Leffler function.

Lemma 1 *For all* $\alpha>0,\beta>0$ *the following relations hold*

$$z^2E_{\alpha,\beta+2\alpha}(z)=E_{\alpha,\beta}(z)-\frac{1}{\Gamma(\beta)}-\frac{z}{\Gamma(\beta+\alpha)}\qquad(18)$$

$$z^3E_{\alpha,\beta+3\alpha}(z)=E_{\alpha,\beta}(z)-\frac{1}{\Gamma(\beta)}-\frac{z}{\Gamma(\beta+\alpha)}-\frac{z^2}{\Gamma(\beta+2\alpha)}\qquad(19)$$

$$z^4E_{\alpha,\beta+4\alpha}(z)=E_{\alpha,\beta}(z)-\frac{1}{\Gamma(\beta)}-\frac{z}{\Gamma(\beta+\alpha)}-\frac{z^2}{\Gamma(\beta+2\alpha)}-\frac{z^3}{\Gamma(\beta+3\alpha)}\qquad(20)$$

◁Let us prove one of these relations:

$$z^2E_{\alpha,\beta+2\alpha}(z)=\sum_{k=0}^{\infty}\frac{z^{k+2}}{\Gamma(\alpha k+\beta+2\alpha)}=\begin{Bmatrix}k+2=n\\k=0\to n=2\\k\to\infty\to n\to\infty\end{Bmatrix}$$

$$=\sum_{n=2}^{\infty}\frac{z^n}{\Gamma(\alpha n+\beta)}$$

For returning the form of two-parametric Mittag-Leffler function, we need initial two elements in the last series, so we will change it a bit

$$z^2E_{\alpha,\beta+2\alpha}(z)=\sum_{n=0}^{\infty}\frac{z^n}{\Gamma(\alpha n+\beta)}-\frac{1}{\Gamma(\beta)}-\frac{z}{\Gamma(\alpha+\beta)}.$$

3. Integral Relations

Using the well-known discrete orthogonality relation

$$\sum_{h=0}^{m-1} e^{i2\pi hk/m} = \begin{cases} m, & \text{if } k \equiv 0 \pmod m \\ 0, & \text{if } k \not\equiv 0 \pmod m \end{cases}$$

And definition (1) of the function $E_{\alpha.\beta}(z)$ we have

$$\sum_{h=0}^{m-1} E_{\alpha,\beta}\left(ze^{i2\pi hk/m}\right) = mE_{\alpha/m,\beta}\left(z^m\right) \quad (m \geq 1) \tag{21}$$

Substituting here $m\alpha$ for α and $z^{1/m}$ for z we obtain

$$E_{\alpha,\beta}(z) = \frac{1}{m}\sum_{h=0}^{m-1} E_{m\alpha,\beta}\left(z^{1/m}e^{i2\pi h/m}\right) \quad (m \geq 1). \tag{22}$$

Similarly, the formula

$$E_{\alpha,\beta}(z) = \frac{1}{2m+1}\sum_{h=-m}^{m} E_{(2m+1)\alpha,\beta}\left(z^{1/(2m+1)}e^{i2\pi h/(2m+1)}\right) \quad (m \geq 0) \tag{23}$$

can be obtained via the relation

$$\sum_{h=-m}^{m} ee^{i2\pi hk/(2m+1)} = \begin{cases} 2m+1, & \text{if } k \equiv 0 \pmod{2m+1}, \\ 0, & \text{if } k \not\equiv 0 \pmod{2m+1}. \end{cases}$$

Using (1) and term-by-term integration we arrive at

$$\int_0^z E_{\alpha,\beta}\left(\lambda t^\alpha\right)t^{\beta-1}dt = z^\beta E_{\alpha,\beta+1}\left(\lambda z^\alpha\right) \quad (\beta > 0) \tag{24}$$

And furthermore, at the more general relation

$$\frac{1}{\Gamma(\alpha)}\int_0^z (z-t)^{\mu-1} E_{\alpha,\beta}\left(\lambda t^\alpha\right)t^{\beta-1}dt = z^{\mu+\beta-1}E_{\alpha,\mu+\beta}\left(\lambda z^\alpha\right) \quad (\mu > 0, \beta > 0) \tag{25}$$

Where the integration is performed along the straight line connecting the points 0 and z.

It follows from formulas (25), (22) and (23) that

$$\frac{1}{\Gamma(\beta)}\int_0^z (z-t)^{\beta-1} e^{\lambda t}dt = z^\beta E_{1,\beta+1}(\lambda z) \quad (\beta > 0) \tag{26}$$

$$\frac{1}{\Gamma(\beta)}\int_0^z (z-t)^{\beta-1} \cosh\sqrt{\lambda}t\,dt = z^\beta E_{2,\beta+1}\left(\lambda z^2\right) \quad (\beta > 0) \tag{27}$$

$$\frac{1}{\Gamma(\beta)}\int_0^z (z-t)^{\beta-1} \frac{\sinh\sqrt{\lambda}t}{\sqrt{\lambda}}dt = z^{\beta+1}E_{2,\beta+2}\left(\lambda z^2\right) \quad (\beta > 0). \tag{28}$$

Here we use series representations of the functions, $e^{\lambda t}$, $\cosh\sqrt{\lambda}t$, $\sinh\sqrt{\lambda}t$ in order to evaluate the integrals above.

◁ Let us prove the relation

$$z^{\beta-1}E_{\alpha,\beta}\left(z^{\alpha}\right)=z^{\beta-1}E_{2\alpha,\beta}\left(z^{2\alpha}\right)+ \\ +\frac{1}{\Gamma(\alpha)}\int_0^z (z-t)^{\alpha-1}E_{2\alpha,\beta}\left(t^{2\alpha}\right)t^{\beta-1}dt \quad (\beta>0) \tag{29}$$

First of all, we have by direct evaluations

$$\int_0^z E_{2\alpha,\beta}\left(t^{2\alpha}\right)t^{\beta-1}\left\{1+\frac{(z-t)^{\alpha}}{\Gamma(\alpha+1)}\right\}dt=$$

$$=\sum_{k=0}^{\infty}\frac{1}{\Gamma(2k\alpha+\beta)}\int_0^z t^{2k\alpha+\beta-1}\left\{1+\frac{(z-t)^{\alpha}}{\Gamma(\alpha+1)}\right\}dt$$

$$=z^{\beta}\sum_{k=0}^{\infty}\frac{z^{2k\alpha}}{\Gamma(2k\alpha+\beta+1)}+z^{\beta}\sum_{k=0}^{\infty}\frac{z^{(2k+1)\alpha}}{\Gamma((2k+1)\alpha+\beta+1)}$$

$$=z^{\beta}\sum_{k=0}^{\infty}\frac{z^{k\alpha}}{\Gamma(k\alpha+\beta+1)}=z^{\beta}E_{\alpha,\beta+1}\left(z^{\alpha}\right)$$

This relations and formula (24) imply

$$\int_0^z E_{2\alpha,\beta}\left(t^{2\alpha}\right)t^{\beta-1}\left\{1+\frac{(z-t)^{\alpha}}{\Gamma(\alpha+1)}\right\}dt$$

$$=\int_0^z E_{\alpha,\beta}\left(t^{\alpha}\right)t^{\beta+1}dt \quad (\beta>0)$$

Differentiating of this formula with respect to z gives us formula (29).

◁ Let us prove the formula

$$\int_0^l z^{\beta-1}E_{\alpha,\beta}\left(\lambda x^{\alpha}\right)(l-x)^{\nu-1}E_{\alpha,\nu}\left(\lambda^{*}(l-x)^{\alpha}\right)dx \\ =\frac{\lambda E_{\alpha,\beta+\nu}\left(l^{\alpha}\lambda\right)-\lambda^{*}E_{\alpha,\beta+\nu}\left(l^{\alpha}\lambda^{*}\right)}{\lambda-\lambda^{*}}l^{\beta+\nu-1} \quad (\beta>,\nu>0) \tag{30}$$

Where λ and $\lambda^{*}\left(\lambda\neq\lambda^{*}\right)$ are any complex parameters.

Indeed, using (1) for any λ, $\lambda^{*}\left(\lambda\neq\lambda^{*}\right)$ and $\beta>0, \nu>0$ we find

$$\int_0^l x^{\beta-1}E_{\alpha,\beta}(\lambda x^\alpha)(l-x)^{\nu-1}E_{\alpha,\nu}(\lambda^*(l-x)^\alpha)\,dx$$
$$=\sum_{n=0}^{\infty}\sum_{m=0}^{\infty}\frac{\lambda^n(\lambda^*)^m}{\Gamma(n\alpha+\beta)\,\Gamma(m\alpha+\nu)}\int_0^l x^{n\alpha+\beta-1}(l-x)^{m\alpha+\nu-1}\,dx$$
$$=\sum_{n=0}^{\infty}\sum_{m=0}^{\infty}\frac{\lambda^n(\lambda^*)^m l^{(n+m)\alpha+\beta+\nu-1}}{\Gamma((m+n)\alpha)+\beta+\nu)}=l^{\beta+\nu-1}\sum_{n=0}^{\infty}\sum_{k=n}^{\infty}\frac{\lambda^n(\lambda^*)^{k-n}\,l^{k\alpha}}{\Gamma(k\alpha+\beta+\nu)}$$
$$=l^{\beta+\nu-1}\sum_{k=0}^{\infty}\frac{(\lambda^*)^k\,l^{k\alpha}}{\Gamma(k\alpha+\beta+\nu)}\sum_{n=0}^{k}\left(\frac{\lambda}{\lambda^*}\right)^n=\frac{l^{\beta+\nu-1}}{\lambda-\lambda^*}\sum_{k=0}^{\infty}\frac{l^{k\alpha}(\lambda^{k+1}-(\lambda^*)^{k+1})}{\Gamma(k\alpha+\beta+\nu)}.$$

Using formula (1) once more, we arrive at (30).

Finally, we obtain two integral relations:

$$\int_0^{+\infty}\mathrm{e}^{-t}E_{\alpha,\beta}(zt^\alpha)t^{\beta-1}\mathrm{d}t=\frac{1}{1-z}\quad(\beta>0,\ |z|<1),\tag{31}$$

$$\int_0^{+\infty}\mathrm{e}^{-t^2/(4x)}\,E_{\alpha,\beta}(t^\alpha)\,t^{\beta-1}\,\mathrm{d}t=\sqrt{\pi}x^{\beta/2}\,E_{2\alpha,\frac{1+\beta}{2}}\left(x^{2\alpha}\right)\ (\beta>0,\ x>0).\tag{32}$$

First of all, since the Mittag-Leffler function (1) is an entire function of order $\rho=1/(\mathrm{Re}\,\alpha)$ and type $\sigma=1$, we have for any $\sigma>1$ the estimate:

$$\left|E_{\alpha,\beta}(z)\right|\le Ce^{\sigma|z|^p},\ |z|\ge|z_\sigma|.$$

Consequently, the integrals in the formulas (31) and (32) are convergent.

Mittag-Leffler type function $E_{\alpha,\alpha}(z)$ plays an essential role in the linear Abel integral equation of the second kind.

Theorem 1. *Let a function $f(t)$ be in the function space $L_1(0,l)$. Let $\alpha>0$ and λ be an arbitrary complex parameter. Then the integral equation*

$$u(t)=f(t)+\frac{\lambda}{\Gamma(\alpha)}\int_0^t(t-\tau)^{\alpha-1}u(\tau)d\tau,\ t\in(0,l)\tag{33}$$

has a unique solution

$$u(t)=f(t)+\lambda\int_0^t(t-\tau)^{\alpha-1}E_{\alpha,\alpha}\left[\lambda(t-\tau)^\alpha\right]f(\tau)d\tau,\ t\in(0,l)\tag{34}$$

in the space $L_1(0,l)$.

This result was discovered in the pioneering work of Hille and Tamarkin.

◁ *Proof.* According to a course of Integral Equations, this is a particular form of Volterra integral equation. In our case, the kernel of the given integral equation is equal to $K(t,\tau)=\frac{(t-\tau)^{\alpha-1}}{\Gamma(\alpha)}$. In order to find the solution of this integral equation, we can use iterative kernels, which are denoted below:

$$K_1(t,\tau)=K(t,\tau)$$

$$K_{n+1}(t,\tau)=\int_\tau^t K(t,z)K_n(z,\tau)dz \quad (n=1,2,...)$$

Then we can easily find the solution with the means of resolvent of this integral equation,

$$u(t)=f(t)+\lambda\int_0^t R(t,\tau;\lambda)f(\tau)d\tau$$

The resolvent is denoted

$$R(t,\tau;\lambda)=\sum_{v=0}^{\infty}\lambda^v K_{v+1}(t,\tau)$$

In this case, it is not easy to notice that the resolvent is equal to

$$R(t,\tau;\lambda)=(t-\tau)^{\alpha-1}E_{\alpha,\alpha}\left[\lambda(t-\tau)^{\alpha}\right]$$

4. Conclusion

The two-parameteric Mittag-Leffler function $E_{\alpha,\beta}$ has a fundamental importance in fractional calculus, and it appears frequently in the solutions of fractional differential and integral equations. However, the expense of calculating this function often prompts efforts to devise accurate approximations that are more cost-effective.

REFERENCES

1. Gorenflo R., Kilbas A. A., Rogosin A. V. On the generalized Mittag-Leffler type function. Integral Transforms Spec. Func., Springer, Germany, 1998.

2. Gorenflo R., Kilbas A. A., Mainardi F., Rogosin S. Mittag-Leffler Functions, Related Topics and Applications; Springer Monographs in Mathematics, Springer Verlag, Berlin, Heidelberg, 2014.

3. Erdeyli A., Magnus W., Oberhettinger F., Tricomi F.G.: Higher Transcendental Functions, vol. 1. McGrew Hill, New York (1953)

4. Salim T.O., Faraj A.W. A generalization of Mittag-Leffler function and integral operator associated with fractional calculus. J Frac. Calc. Appl. 2012

5. M. M. Dzzherbashian [Djrbashian], Integral Transforms and Representation of Functions in the Complex Domain, Nauka, Moskow, 1996 (Russian).

6. S. Gerhold, Asmptotics for a variant of the Mittag-Leffler function, Integral Transforms Spec. Funct., 2012

7. Samko S G, Kilbas AA, Marichev OI. Fractional integrals and derivatives: Theory and Applications.

8. A. A. Kilbas and M. Saigo, H-Transform. Theory and Applications, Chapman and Hall/CRC, Boca Raton, London, New York, Washington, D. C., 2004.

9. A. M. Mathai, Mittag-Leffler function, Ch. 2, in A.M. Mathai and H.J. Haubold (eds.), An Introduction to Fractional Calculus, Nova Science Publishers, 2017.

10. R. N. Pillai, On Mittag-Leffler functions and related distributions, Ann. Inst. Stat. Math., 1990.

11. A. K. Shukla and J.C. Prajapati, On a generalization of Mittag-Leffler function and its properties, J. Math. Anal. Appl.,2007.

M U N D A R I J A

№		Bet
1	IKKI PARAMETRLI MITTAG–LEFFLER FUNKSIYASINING ASOSIY XOSSALARI VA KASR TARTIBLI DIFFERENSIAL TENGLAMALARDAGI QO‘LLANILISHI ***Ne'matjonov Furqatjon Farhodjon o‘g‘li***	**6**
2	THE TWO-PARAMETRIC MITTAG-LEFFLER FUNCTION AND ITS SOME PROPERTIES ***Furkatjon F. Nematjonov***	**8**
3	KO‘P KOMPONENTLI KOLLOID SUYUQLIK TOMCHISINING BUG‘LANISHIDA KONVEKSIYA-DIFFUZIYA VA KAPILLYAR OQIMLAR FIZIKASI ***Madaminov Ikromjon Baxtiyor o‘g‘li***	**16**
4	GENERATIV AI YORDAMIDA TEXNIK HUJJATLARNI AVTOMATIK YARATISH TEXNOLOGIYALARI ***Mamirov Diyorbek Rasiljon o‘g‘li***	**22**
5	KOLLOID SUYUQLIK TOMCHISINING KLINOVIDLI DEGIDRATATSIYASIDA FAZAVIY O‘TISH VA STRUKTURA HOSIL BO‘LISHINING FIZIK MEXANIZMLARI ***Madaminov Ikromjon Baxtiyor o‘g‘li***	**31**
6	SnO_2 PLYONKALARINING TURLI GAZLARINI SEZUVCHANLIK XOSSALARIGA SIRTIY NUQSONLARNI TA'SIRI ***Toxirova Mashhuraxon***	**37**
7	HARORAT MAYDONIDA DISTILLANGAN SUV VA NaCl ERITMASI TOMCHILARINING GEOMETRIK EVOLYUTSIYASI VA EVAPORATSIYA KINETIKASI ***Madaminov Ikromjon Baxtiyor o‘g‘li***	**41**
8	KONTAKT BURCHAGI EVOLYUTSIYASI ORQALI KOLLOID TOMCHILARNING CCR, CCA VA ARALASH QURISH REJIMLARINI FIZIK MODELLASHTIRISH ***Madaminov Ikromjon Baxtiyor o‘g‘li***	**44**
9	О СУЩЕСТВОВАНИИ РЕШЕНИЯ ЗАДАЧИ ТИПА ДИРИХЛЕ ДЛЯ ЭЛЛИПТИЧЕСКОГО УРАВНЕНИЯ С ТРЕМЯ СИНГУЛЯРНЫМИ КОЭФФИЦИЕНТАМИ	**47**